



Original Article

Socio-Demographic Risk Factors for Severe Acute Malnutrition

Naznin Akther¹, Md. Tarek Azad², Tahmina Jahan Chowdhury³, Md. Muazzem Hussain⁴,
Forida Akter⁵, Aloka Nanda Talukder⁶

¹Assistant Professor, Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College, Sylhet.

²Professor, Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College, Sylhet.

^{3,4}Associate Professor, Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College, Sylhet.

^{5,6}Indoor Medical Officer, Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College, Sylhet.

ABSTRACT

Malnutrition is one of the leading causes of morbidity and mortality in under-five children in developing countries. Bangladesh being one of the developing countries, malnutrition is an important public health problem. This was a hospital-based case-control study conducted in the department of Paediatrics, Jalalabad Ragib-Rabeya Medical College Hospital, Sylhet during the period from July 2013 to June 2014 to see the socio-demographic risk factors for severe acute malnutrition among children under five. Forty five severe acute malnourished children, aged 6 months to 5 years, with a weight-for-height Z score below -3 were enrolled in the case group. Controls were age and sex matched, well nourished children with a weight-for-height Z score of -1 to <+2 who were admitted at the same time following the admission of the case due to other problems. Rural participants ($p<0.001$), undernourished mother ($p=0.006$), less-educated parents ($p<0.001$), lower monthly income ($p=0.005$), birth order 3 or more ($p=0.017$), family member 5 or higher ($p=0.015$), lack of exclusive breast feeding ($p<0.001$), and inappropriate weaning ($p<0.001$) significantly increased the risk of development of severe acute malnutrition. To reduce childhood malnutrition, due emphasis should be given to improve the socioeconomic status, parental education, knowledge and practice of parents on appropriate infant and child feeding practices.

Keywords: Severe acute malnutrition, Socio-demographic risk factors, Under-five children.

[Jalalabad Med J 2022; 19 (2): 47-53]

INTRODUCTION

Severe malnutrition among children, and particularly acute malnutrition, is a life-threatening condition that requires urgent intervention. The condition can also be chronic. The World Health Organization (WHO) defines severe acute malnutrition (SAM) as 'a very low weight for height, by visible severe wasting, or by the presence of nutritional oedema'¹. Wasting is defined in children aged 6-59 months as a weight for height Z score below

-3 standard deviations (SD) of the WHO growth standards and/or a mid-upper arm circumference (MUAC) <115 mm².

Severe acute malnutrition affects an estimated 20 million children under the age of five worldwide and needs treatment urgently. If it is untreated, the median under-five case-fatality rate for SAM ranges from 30% to 50%. SAM kills one million children each year worldwide, an average of one child every thirty seconds^{1,3,4,5}. Malnutrition alone is one of the leading causes of about 10% (140 million disability adjusted life-years) of the global burden of disease⁶. To reduce

Address of Correspondence:

Dr. Naznin Akther, Assistant Professor, Department of Paediatrics,
Jalalabad Ragib-Rabeya Medical College, Sylhet. Mobile:
01717414470, Email: drnazninakter27@gmail.com.

the mortality rate, community and facility-based health services are thus principally focused on identifying and treating children with SAM⁷. Improving SAM survival is a global public health priority and key to the achievement of international health targets such as Millennium Development Goal 4, reducing child mortality⁸.

Bangladesh has the world's fourth highest number of children suffering from SAM⁹. In Bangladesh, the proportion of children with severe wasting is 2.9%, thus the total number being 500,000¹⁰. Different studies have shown that several factors closely related to infant and child malnutrition, including mother's nutritional status, child's sex, family size, low dietary intake, breast feeding status, poverty, socio-economic status, place of residence, child care behaviour, previous birth interval, early and delayed weaning, cultural beliefs, access to health care, and environmental ecosystems^{11,12,13}.

Although factors predisposing children to protein energy malnutrition have been studied at home¹⁴ and abroad^{15,16,17}, there is a lack of study on socio-demographic risk factors amongst under five children with malnutrition in Sylhet. So, this study was designed to find out socio-demographic risk factors for severe acute malnutrition amongst children attending Jalalabad Ragib-Rabeya Medical College Hospital, Sylhet.

MATERIALS AND METHODS

This was a hospital-based case-control study conducted in the Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College Hospital, Sylhet during the period from July 2013 to June 2014. Children with SAM aged between 6 months and 5 years with a weight for height Z score of <-3 , admitted to the Department of Paediatrics, Jalalabad Ragib-Rabeya Medical College Hospital, Sylhet were evaluated. Those who fulfilled the inclusion and exclusion criteria were enrolled in the case group. In this way, forty-five severe acute malnourished children aged 6 months to 5 years with a weight for height Z score of <-3 were consecutively enrolled in this study. Children with chronic malnutrition, the presence of any congenital anomaly or chronic disease, and those who refused to be enrolled in this study were excluded. Forty five cases were matched with ninety well-nourished controls of similar age and sex with weight for height Z score -1 to $+2$, who were admitted at the same time following the admission of the case due to other problems (Case-control ratio 1:2). The aims and

objectives of the study were explained to the mother or the attendant of the children in an easily understandable local language, and then written informed consent was taken from the mother or the attendant. Mothers of the selected children were interviewed to obtain information on their socioeconomic and demographic issues through pretested structured questionnaires. This included maternal age, parental education, monthly family income, occupation, and birth order. The children's information included their age, gender, breast feeding status, weaning practice, place of residence, and number of family members. The weight of the child was measured to the nearest 0.1 kg and their height was measured to the nearest 0.1 centimetre. For children aged less than two years, weight was recorded using a conventional beam balance scale. Recumbent length and standing height of under five children were measured with an infantometer and stadiometer accordingly. The age of the children was recorded either from the mother's statement or from the discharge paper of the mother after delivery. The severe acute malnutrition was diagnosed according to the child growth standard of the World Health Organization (WHO). The Anthro 2005 software of the WHO was used for calculating the Z-score. The same questionnaire was applied to the control group, which consisted of children without malnutrition admitted for other medical problems. Data was processed and analyzed with the help of the computer program Statistical Package for Social Sciences (SPSS) version 16. Quantitative data was expressed as mean and standard deviation, and qualitative data as frequency and percentage. Comparison was done by the Chi-square (χ^2) test and the Z-test where applicable. Socio-demographic risk factors were calculated using an Odd ratio with a confidence interval. A probability (p) value of <0.05 was considered statistically significant.

RESULT

In our study, 135 children were studied, of whom 45 were severely malnourished from age 6 months to 5 years, and 90 were age and sex matched well-nourished controls. The mean age was 17.33 ± 8.2 months in the case group and 16.48 ± 7.28 months in the control group. In the case group, 25 (55.6%) were male and 20 (44.4%) were female, whereas in the control group, 55 (61.1%) were male and 35 (38.9%) were female. Age and sex didn't differ significantly between the case and control groups. (Table-I, Figure-1).

Table-I: Distribution of the case and control by age at admission, N=135

Age (Months)	Case group, n (%)	Control group, n (%)	p-value
6-12	15 (33.3)	31 (34.4)	*0.720
13-24	25 (55.6)	53 (58.9)	
25-36	5 (11.1)	6 (6.7)	
Mean age (Months±SD)	17.33±8.2	16.48±7.28	†0.322

*Chi-square (χ^2) test and †Z test were employed to analyze the data.




Figure-1: Distribution of case and control by gender, N=135

Table-II: Socio-demographic risk factors of case and control, N=135

Risk Factors		Case n (%)	Control n (%)	p-value	Odds ratio (95% of CI)
Place of residence	Urban	12 (26.7)	58 (64.4)	<0.001	4.984 (2.264-10.974)
	Rural	33 (73.3)	32 (35.6)		
Breast feeding status	Exclusive	12 (26.7)	61 (67.8)	<0.001	0.173 (0.078-0.383)
	Non-exclusive	33 (73.3)	29 (32.2)		
Weaning	Appropriate	8 (17.8)	53 (58.9)	<0.001	0.151 (0.063-0.361)
	Inappropriate	37 (82.2)	37 (41.1)		
Number of family member	> 5	40 (88.9)	63 (70)	0.015	3.429 (1.220-9.635)
	< 5	5 (11.1)	27 (30)		

The case group had more children living in rural areas (73.3%) than the control group (35.6%), which was statistically significant (OR=4.984, 95% CI=2.264-10.974, $p=0.001$). Exclusive breast feeding was significantly lower in the case group (26.7%) than in the control group (67.8%) (OR=0.173; 95% of CI=0.078-0.383; $p<0.001$). Appropriate weaning practice was lower in the case group (17.8%) than in the control group (58.9%), which was statistically

Only 8.9% of mothers had more than 5 years of schooling in the case group, which was 72% in the control group. (OR=28.188; 95% of CI=9.124-87.082; $p<0.001$). Working mothers made up 4.4% of the case group and 8.9% of the control group among the mothers, however there was no statistically significant difference between the two (OR=2.098; 95% CI=0.427-10.314; $p=0.353$). In the case group, 42.2% of fathers had more than 5 years of schooling, which

Table -III: Socio-demographic risk factors of parents of case and control, N=135

Risk Factors		Case n (%)	Control n (%)	p-value	Odds ratio (95% of CI)
Mean age of the mother (Years±SD)		25.13±4.97	25.78±3.25	†0.162	2.00 (0.749-5.343)
Mother's BMI (kg/m ²)	<18.5	12 (26.7)	8 (8.9)	*0.006	3.727
	> 18.5	33 (73.3)	82 (91.1)		(1.397-9.947)
Mother's education	<5 yrs of schooling	41 (91.1)	24 (26.7)	*<0.00	28.188
	>5 yrs of schooling	4 (8.9)	66 (73.3)	1	(9.124-87.082)
Mother's occupation	Housewife	43 (95.6)	82 (91.1)	*0.353	2.098
	Service	2 (4.4)	8 (8.9)		(0.427-10.314)
Father's education	<5 yrs of schooling	26 (57.8)	11 (24.4)	*<0.00	4.230
	>5 yrs of schooling	19 (42.2)	34 (75.6)	1	(1.974-9.065)
Monthly income	<5000 taka	13 (28.9)	9 (10)	*0.005	3.656
	>5000 taka	32 (71.1)	81 (90)		(1.424-9.392)
Birth order	> 3	12 (26.7)	7 (7.8)	*0.017	3.388
	<3	33 (73.3)	83 (92.2)		(1.193-9.619)

*Chi-square (χ^2) test and †Z test were employed to analyze the data.

significant (OR=0.151; 95% of CI=0.063-0.361; $p<0.001$). The numbers of family members were greater in the case group than in the control group. (OR=3.429; 95% of CI=1.220-9.635; $p=0.015$) (Table-II).

The mean age of the mother of both cases and controls showed no significant difference (OR=2.00; 95% of CI=0.749-5.343; $p=0.162$). But the mothers' BMI were lower in the case group than in the control group (OR=3.727; 95% of CI=1.397-9.947; $p=0.006$).

was 75.6% in the control group. The difference was statistically significant (OR=4.230; 95% of CI=1.974-9.065; $p<0.001$). Approximately thirty percent (28.9%) of the children in the case group came from low-income families, whereas only 10% of the children in the control group came from low-income families (OR=3.656; 95% of CI=1.424-9.392; $p=0.005$). The number of birth order >3 was higher (26.7%) in the case group than in the control group (7.8%), which was statistically significant (OR=3.388; 95% of CI=1.193-9.619; $p=0.017$) (Table-III).

DISCUSSION

The aim of the study was to determine the socio-demographic risk factors for SAM in a tertiary level hospital, so that we might be able to take appropriate action to reduce the burden of childhood malnutrition.

Among 45 cases and 90 controls the mean age of the cases was 17.33 ± 8.2 months and the mean age of the controls was 16.48 ± 7.28 months. In the present study 25 (55.6%) patients were male and 20 (44.4%) patients were female in case group; whereas 55 (61.1%) subjects were male and 35 (38.9%) subjects were female in the control group. Age and sex didn't differ significantly between case and control groups. This result was consistent with the study of Nahar et al¹⁹. They found that the mean age of the patients with severely-underweight children was 12.6 ± 4 months in case group; whereas it was 16.48 ± 7.28 in the control group. In the case group, 56.6% of patients with severely-underweight were male and 43.6% were female; whereas in control group 56% of subjects were male and 44% were female. There was no significant difference in age and sex between the cases and controls.

In the current study the mean age of mother was 25.13 ± 4.97 years in case group; whereas 25.78 ± 3.25 years in control group. There was no statistically significant difference of the age group of the mothers between case and control group.

Underweight mothers have a greater risk of their children being malnourished. A healthy mother can have healthy children. Children of well nourished mother had a lower risk of being underweight compared to undernourished mother¹¹. In our study maternal BMI less than $<18.5 \text{ kg/m}^2$ (Under nourished mother) significantly increased the risk of development of severe acute malnutrition. This result correlated with the study of Nahar et al¹⁹.

This study showed that the risk of development of severe acute malnutrition was significantly higher in rural participants. Siddiqi et al.¹¹ and Yimer G²⁰ found the similar result in their study. Availability of water and sanitary facilities, better socioeconomic status and child health care facilities in urban areas results into less risk of being malnourished among children living in urban areas than their rural counterparts¹¹.

In a study done by Rayhan and Khan²¹ reported that the risk of malnutrition decreased as paternal education increased. In our study, <5 years of schooling of the parents significantly increased the risk of development

of severe acute malnutrition in children. This result was consistent with the study of Nahar et al¹⁹. Maternal education is known to have profound beneficial effects on child feeding, health seeking and care giving practices which is especially important for child nutritional status¹¹.

Maternal occupation did not differ significantly between cases and controls in this study. This result was in agreement with the study of Dereje N²² who found that there was no significant difference in the child's severe wasting in cases and controls in relation to the maternal occupation ($p > 0.05$). However, the opposite result was found by Nahar et al.¹⁹, where mothers working outside the home were significantly associated with the development of severe acute malnutrition in children.

In this study, lower monthly income (<5000 Taka) and birth order of 3 or more significantly increased the risk of development of severe acute malnutrition. This result was in agreement with the study of Nahar et al¹⁹. In the current study, the number of family members 5 or higher significantly increased the risk of development of severe acute malnutrition. In a study, Dereje N²² found that 86.1% of the households in the cases had more than 5 family members as compared to 76.9% of the controls.

In the present study, exclusive breastfeeding for the first 6 months of life and appropriate weaning practices significantly reduced the risk of the development of severe acute malnutrition. Nahar et al.¹⁹ in their study showed that predominant breastfeeding for <4 months and early weaning significantly increased the risk of the development of severe acute malnutrition, which supports our result.

CONCLUSION

This present study concluded that rural residence, undernourished mother, less-educational qualification of the parents, lower monthly income, higher birth order, large family member, lack of exclusive breastfeeding and inappropriate weaning practice are some socio-demographic risk factors for development of severe acute malnutrition. To reduce the burden of childhood malnutrition, due emphasis should be given to improving the socioeconomic status, parental education, maternal health, and parents' knowledge about appropriate feeding practice for children.

REFERENCES

1. World Health Organization. Severe acute malnutrition. Geneva: World Health Organization; 2013 [Cited 2014 May 12]. Available at: <http://www.concernusa.org/media/pdf/2007/10/CommunityBased%20Management%20of%20Severe%20Acute%20Malnutrition.pdf>.
2. World Health Organization. Management of severe malnutrition: a manual for physicians and other senior health workers. Geneva: World Health Organization; 1999 [Cited 2014 May 12]. Available at: <http://whqlibdoc.who.int/hq/1999/a57361.pdf>.
3. World Health Organization. Innovative approach tackles malnutrition in the community: a joint statement by the WHO, the World Food Programme (WFP), the United Nations Standing Committee on Nutrition (SCN) and UNICEF. Geneva: World Health Organization; 2007 [Cited 2014 May 12]. Available at: <http://www.who.int/mediacentre/news/releases/2007/pr27/en/>.
4. Collins S, Dent N, Binns P, Bahwere P, Sadler K, Hallam A. Management of severe acute malnutrition in children. *Lancet* 2006; 368 (9551): 1992-2000.
5. Fergusson P, Tomkins A, Kerac M. Improving survival of children with severe acute malnutrition in HIV-prevalent settings. *Int Health* 2009; 1 (1): 10-6.
6. Ezzati M, Lopez AD, Rodgers A, Hoorn SV, Murray CJL, Comparative Risk Assessment Collaborating Group. Selected major risk factors and global and regional burden of disease. *Lancet* 2002; 360 (9343): 1347-60.
7. Shams Z, Zachariah R, Enarson DA, Satyanarayana S, Van den Bergh R, Ali E, et al. Severe malnutrition in children presenting to health facilities in an urban slum in Bangladesh. *Public Health Action* 2012; 2 (4): 107-11.
8. United Nations. Millennium Development Goals. New York: United Nations; 2008 [Cited 2013 March 26]. Available at: http://www.un.org/millenniumgoals/2008highlevel/pdf/newsroom/mdg%20reports/MDG_Report_2008_ENGLISH.pdf.
9. Sadler K, Puett C, Mothabbir G, Myatt M. Community case management of severe acute malnutrition in southern Bangladesh. Boston: Freistein International Center; 2011 [Cited 2013 March 26]. Available at: <http://www.scribd.com/doc/69741063/Community-Case-Management-of-Severe-Acute-Malnutrition-in-Southern-Bangladesh>.
10. Institute of Public Health Nutrition, Directorate General of Health Services, Ministry of Health and Family Welfare. Training modules on management of children with severe acute malnutrition in Bangladesh. Dhaka: Institute of Public Health Nutrition, Directorate General of Health Services, Ministry of Health and Family Welfare, Government of the People's Republic of Bangladesh; 2011.
11. Siddiqi MNA, Haque MN, Goni MA. Malnutrition of under-five children: evidence from Bangladesh. *Asian J Med Sci* 2011; 2 (2): 113-9.
12. Kamal SMM. Socio-economic Determinants of severe and moderate stunting among under-five children of rural Bangladesh. *Mal J Nutr* 2011; 17 (1): 105 -18.
13. Ayaya SO, Esamai FO, Rotich J, Olwambula AR. Socio-economic factors predisposing under five-year-old children to severe protein energy malnutrition at the Moi teaching and referral hospital, Eldoret, Kenya. *East Afr Med J* 2004; 81 (8): 415-21.
14. Rahman A, Chowdhury S. Determinants of chronic malnutrition among preschool children in Bangladesh. *J Biosoc Sci* 2007; 39 (2): 161-73.
15. Owor M, Tumwine JK, Kikafunda JK. Socio-economic risk factors for severe protein energy malnutrition among children in Mulago Hospital, Kampala. *East Afr Med J* 2000; 77 (9): 471-5.
16. Roy NC. Use of mid-upper arm circumference for evaluation of nutritional status of children and for identification of high-risk groups for malnutrition in rural Bangladesh. *J Health Popul Nutr* 2000; 18 (3): 171-80.
17. el-Sayed N, Mohamed AG, Nofal L, Mahfouz A, Zeid HA. Malnutrition among pre-school children in Alexandria, Egypt. *J Health Popul Nutr* 2001; 19 (4): 275-80.
18. Jesmin A, Yamamoto SS, Malik AA, Haque MA. Prevalence and determinants of chronic malnutrition among preschool children: a cross-sectional study in

- Dhaka City, Bangladesh. *J Health Popul Nutr* 2011;29 (5): 494-9.
19. Nahar B, Ahmed T, Brown KH, Hossain MI. Risk factors associated with severe underweight among young children reporting to a diarrhoea treatment facility in Bangladesh. *J Health Popul Nutr* 2010; 28 (5): 476-83.
 20. Yimer G. Malnutrition among children in Southern Ethiopia: levels and risk factors. *Ethiopian J Health Dev* 2000; 14 (3): 283-92.
 21. Rayhan MI, Khan MSH. Factors causing malnutrition among under five children in Bangladesh. *Pak J Nutr* 2006; 5 (6): 558-62.
 22. Dereje N. Determinants of severe acute malnutrition among under five children in Shashogo Woreda, Southern Ethiopia: a community based matched case control study. *J Nutr Food Sci* 2014; 4 (5). doi: 10.4172/2155-9600.1000300.